

Abstract Submitted
for the DPP96 Meeting of
The American Physical Society

Sorting Category: 4.6 (experimental)

Further Experimental Investigations of the Richtmyer-Meshkov Instability¹ L.M. LOGORY, P.L. MILLER, T.A. PEYSER, P.E. STRY, Lawrence Livermore National Laboratory — We report on further experimental investigations of the Richtmyer-Meshkov instability from an initially nonlinear perturbation, conducted on the Nova laser. The experiments use a Nova hohlraum as a driver source for a strong shock in a miniature shock tube attached to the hohlraum. The shock tube contains brominated plastic and low-density carbon foam as the two working fluids, with a micro-machined sawtooth interface between them serving as the perturbation. The shock, upon crossing the interface, instigates the Richtmyer-Meshkov instability from the perturbation. The resulting growth of the mixing layer is diagnosed radiographically. We have previously reported upon a results from a single wavelength and amplitude of perturbation ². A study of the effect of variations in amplitude and wavelength on the nonlinear growth of the instability will be discussed.

¹This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48.

²T. A. Peyser *et al.*, *Phys. Rev. Lett.* **75**, 2332 (1996).

☒ Prefer Oral Session
☐ Prefer Poster Session

Paul L. Miller
pmiller@llnl.gov
Lawrence Livermore National Laboratory

Date submitted: July 9, 1996

Electronic form version 1.1